

AMENDMENTS TO THE CLAIMS

The following is a listing of all claims in the application. Claim 1 is amended, Claims 2-5 are canceled, Claims 16-30 are withdrawn, and new Claims 31-45 are added as follows:

1. (currently amended) A bistable molecule for a multiple electrode device comprising at least one pair of electrodes that form at least one junction and at least one ~~connector species~~ said bistable molecule connecting said pair of electrodes in said junction, said junction having a functional dimension in nanometers or micrometers, wherein said ~~“at least one connector species”~~ comprises said bistable molecule provided with includes at least one photosensitive functional group ~~for patterning said connector species~~.

Claims 2-5 (canceled)

6. (original) The bistable molecule of Claim 1 wherein said photosensitive functional group is sensitive to ultraviolet, electron-beam, or X-ray radiation.

7. (original) The bistable molecule of Claim 1 wherein said bistable molecule comprises a main chain and at least one pendant group and wherein at least one photosensitive functional group is attached either to said main chain or to said pendant group.

8. (original) The bistable molecule of Claim 7 wherein one said photosensitive group is attached to at least one end of said bistable molecule.

9. (original) The bistable molecule of Claim 7 wherein said photosensitive group is selected from the group consisting of  $\alpha$ -carboxy-2-nitrobenzyl; 1-(2-nitrophenyl)ethyl; 4,5-dimethoxy-2-nitrobenzyl; 1-(4,5-dimethoxy-2-nitrophenyl)ethyl; (4,5-dimethoxy-2-nitrobenzyloxy)carbonyl; 5-carboxymethoxy-2-nitrobenzyl; [(5-carboxymethoxy-2-nitrobenzyl)oxy]carbonyl; desoxybenzoinyl; and anthraquinon-2-ylmethoxycarbonyl.

10. (original) The bistable molecule of Claim 1 wherein said molecule evidences switching based on electric (E) field induced band gap change, selected from the group consisting of:

(1) an electric field ("E-field") induced rotation of at least one rotatable section (rotor) of a molecule to change the band gap of the molecule (rotor/stator configuration);


(2) E-field-induced charge separation or recombination of the molecule via chemical bonding change to alter the band gap:

(2a) E-field-induced band gap change caused by the change of extended conjugation via charge separation or recombination accompanied by increasing or decreasing  $\pi$ - and/or p-electron localization;

(2b) E-field-induced band gap change caused by a change of extended conjugation via charge separation or recombination and  $\pi$ -bond breaking or formation; and

(3) E-field-induced band gap change via molecular folding or stretching.

11. (original) The bistable molecule of Claim 10 wherein said bistable molecule comprises:



5 where:

$\text{A}^-$  is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its deriva-

tives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

$D^+$  is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH; (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P, wherein said Donor group is more electropositive than said Acceptor group;

$Con_1$  and  $Con_2$  are connecting units between one molecule and another molecule or between a molecule and a substrate, said connecting units containing an attaching unit and at least one of said connecting units containing said photosensitive group, wherein said photosensitive group is selected from the group consisting of photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive heteroring systems with at least one hetero-atom selected from the group consisting of N, O, S, B, and P,, and wherein the attaching unit is selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) hetero atoms selected from the group consisting of N, O, S, B, Se, and P, (e) functional groups with at least one of said hetero atoms (f) hydrocarbons, and (g) substituted hydrocarbons;

$X_1$ ,  $X_2$ ,  $X_3$  are tuning units built into the ring system which serve to tune the electronic properties, the optical properties, or both, of the bistable molecule as well those of the ring system undergo a smooth and desired tautomerization transition under the influence of an applied external E-field, wherein the tuning units are selected from the group consisting of a hetero atom selected from the group consisting of N, P, and As; hydrocarbons; and substituted hydrocarbons;


$G_1$  and  $G_2$  are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more fragments to achieve desired molecular properties, wherein the bridging groups are either (a) selected from the group consisting of (i) hetero atoms selected from the group consisting of N, O, S, and P; (ii) functional groups with at least one of said hetero atoms; (iii) saturated or unsaturated hydrocarbons; and (iv) substituted hy-

drocarbons or (b) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

Q is a connecting unit between two phenyl rings, selected from the group consisting of S, O, NH, NR, hydrocarbons, and substituted hydrocarbons; and

45 H is a hydrogen atom.

12. (original) The bistable molecule of Claim 10 wherein said bistable molecule comprises:



5 where:

A<sup>-</sup> is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

D<sup>+</sup> is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P, wherein said Donor group is more electropositive than said Acceptor group;

Con<sub>2</sub> is a connecting unit between one molecule and another molecule or between a molecule and a substrate, said connecting unit containing an attaching unit and said photosensitive group, wherein said photosensitive group is selected from the group consisting of photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive hetero-ring systems with at least one hetero-atom selected from the group consisting of N, O, S, B, and P and wherein the attaching unit is selected from the group consisting of carboxylic acid and its derivatives; sulfuric acid and its derivatives; phosphoric acid and its derivatives; hetero atoms selected from the group consisting of N, O, S, B, Se, and P functional groups with at least one of said hetero atoms; hydrocarbons; and substituted hydrocarbons;

X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub> are tuning units built into the ring system which serve to tune the electronic properties, the optical properties, or both, of the bistable molecule as well as to that the ring system undergoes a smooth and desired tautomerization transition under the influence of an applied external E-field, wherein the tuning units are selected from the group consisting of a hetero atom selected from the group consisting of N, P, and As; hydrocarbons; and substituted hydrocarbons;


G<sub>1</sub> and G<sub>2</sub> are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more fragments to achieve desired molecular properties, wherein the bridging groups are either (a) selected from the group consisting of (i) hetero atoms selected from the group consisting of N, O, S, and P; (ii) functional groups with at least

one of said hetero atoms; (iii) saturated or unsaturated hydrocarbons; and (iv) substituted hydrocarbons or (b) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

Q is a connecting unit between two phenyl rings, selected from the group consisting of S, O, NH, NR, hydrocarbons, and substituted hydrocarbons; and

H is a hydrogen atom.

13. (original) The bistable molecule of Claim 10 wherein said bistable molecule comprises:



where:

5           A<sup>-</sup> is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

10           D<sup>+</sup> is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group  
15 is more electropositive than said Acceptor group;


          G<sub>1</sub>=G<sub>2</sub>, G<sub>3</sub>=G<sub>4</sub>, G<sub>5</sub>=G<sub>6</sub>, and G<sub>7</sub>=G<sub>8</sub> are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more conjugated rings to achieve desired electronic properties, wherein the bridging groups are either (a) photosensitive functional groups or (b) selected from the group consisting of (i) hetero atoms selected from the  
20 group consisting of N, O, S, and P; (ii) functional groups with at least one of said hetero atoms; (iii) saturated or unsaturated hydrocarbons; and (iv) substituted hydrocarbons, or (c) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

          Con<sub>1</sub> and Con<sub>2</sub> are connecting units between one molecule and another molecule or  
25 between a molecule and a substrate, said connecting units containing an attaching unit and at least one of said connecting units containing said photosensitive group, wherein said photosensitive group is selected from the group consisting of photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive  
30 thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive heteroring systems with at least one hetero-atom selected from the group consisting of N, O, S, B, and P and wherein the attaching unit is selected from the group consisting of carboxylic acid and its derivatives; sulfuric acid and its derivatives; phosphoric acid and its derivatives; hetero atoms selected from the group consisting of N, O, S, B, Se, and P; functional groups with  
35 at least one of said hetero atoms; hydrocarbons, and substituted hydrocarbons;

$R_1$ ,  $R_2$ , and  $R_3$  are spacing groups selected from the group consisting of (a) hydrogen, (b) saturated or unsaturated hydrocarbons, and (c) substituted hydrocarbons; and

$J_1$  and  $J_2$  are tuning groups to provide at least one appropriate functional effect selected from the group consisting of inductive effects, resonance effects, and steric effects, where said tuning groups are selected from the group consisting of (a) hydrogen, (b) hetero atoms selected from the group consisting of N, O, S, P, B, F, Cl, Br, and I, (c) functional groups with at least one of said hetero atoms, (d) saturated or unsaturated hydrocarbons, and (e) substituted hydrocarbons.

14. (original) The bistable molecule of Claim 10 wherein said bistable molecule comprises:



5 where:


$A^-$  is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

$D^+$  is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group is more electropositive than said Acceptor group;

$R_1$ ,  $R_2$ , and  $R_3$  are spacing groups selected from the group consisting of (a) hydrogen, (b) saturated or unsaturated hydrocarbons, and (c) substituted hydrocarbons; and

$J_1$  and  $J_2$  are tuning groups to provide at least one appropriate functional effect selected from the group consisting of inductive effects, resonance effects, and steric effects, said tuning groups are selected from the group consisting of (a) hydrogen, (b) hetero atoms selected from the group consisting of N, O, S, P, B, F, Cl, Br, and I, (c) functional groups with at least one of said hetero atoms, (d) saturated or unsaturated hydrocarbons, and (e) substituted hydrocarbons.

15. (currently amended) The bistable molecule of Claim 10 wherein said bistable molecule comprises:



5 where:

$A^-$  is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its deriva-

tives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

$D^+$  is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group is more electropositive than said Acceptor group;

$G_1=G_2$ ,  $G_3=G_4$ ,  $G_5=G_6$ , and  $G_7=G_8$  are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more conjugated rings to achieve desired electronic properties, wherein the bridging groups are either (a) photosensitive functional groups, or (b) selected from the group consisting of (i) hetero atoms selected from the group consisting of N, O, S, and P; (ii) functional groups with at least one of said hetero atoms; (iii) saturated or unsaturated hydrocarbons; and (iv) substituted hydrocarbons, or (c) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

$Con_1$  and  $Con_2$  are connecting units between one molecule and another molecule or between a molecule and a substrate, said connecting units containing an attaching unit and at least one of said connecting units containing said photosensitive group, wherein said photosensitive group is selected from the group consisting of photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive heteroring ~~system(s?)~~ systems with at least one hetero-atom selected from the group consisting of N, O, S, B, and P and wherein the attaching unit is selected from the group consisting of carboxylic acid and its derivatives; sulfuric acid and its derivatives; phosphoric acid and its derivatives; hetero atoms selected from the group consisting of N, O, S, B, Se, and P; functional groups with at least one of said hetero atoms; hydrocarbons; and substituted hydrocarbons; and

$J_1$ ,  $J_2$ ,  $J_3$ , and  $J_4$  are tuning groups which contain solvent functional groups selected from the group consisting of OH, NHR, COOH, and CN, where R is alkyl or aryl, wherein  $J_1$ -PSG,  $J_2$ -PSG,  $J_3$ -PSG, and  $J_4$ -PSG are linkages of said tuning groups with said photosensitive

groups and are selected from the group consisting of ether, ester, carbonate, amide, and carbamate linkages.

16. (withdrawn) A method for fabricating a multiple electrode device comprising at least one pair of electrodes that form at least one junction and at least one connector species connecting said pair of electrodes in said junction, said junction having a functional dimension in nanometers or micrometers, wherein said at least one connector species comprises said bistable molecule provided with at least one photosensitive functional group for patterning said connector species, said method comprising:

- (a) forming a first set of said electrodes on a substrate;
- (b) depositing a film of said bistable molecule(s) including said photosensitive group prior to said depositing;
- (c) exposing portions of said bistable molecular film to desired radiation; and
- (d) removing unwanted portions of said bistable molecular film to provide a photopatterned molecule.

17. (withdrawn) The method of Claim 16 wherein said at least one pair of electrodes comprises a positive terminal and a negative terminal and wherein said method further comprises, after step (d):

- (e) depositing a second set of said electrodes adjacent said first set of said electrodes.

18. (withdrawn) The method of Claim 17 wherein said second set of electrodes is deposited above said first set of electrodes, at a non-zero angle thereto.

19. (withdrawn) The method of Claim 17 wherein said second set of electrodes is deposited in the same plane as said first set of electrodes.

20. (withdrawn) The method of Claim 17 wherein said at least one pair of electrodes comprises said first set of electrodes and at least one probe addressing tip.

21. (withdrawn) The method of Claim 16 wherein said photosensitive functional group is sensitive to ultraviolet, electron-beam, or X-ray radiation.

22. (withdrawn) The method of Claim 16 wherein said bistable molecule comprises a main chain and at least one pendant group and wherein at least one photosensitive functional group is attached either to said main chain or to said pendant group.

23. (withdrawn) The method of Claim 22 wherein one said photosensitive group is attached to at least one end of said bistable molecule.

24. (withdrawn) The method of Claim 22 wherein said photosensitive group is selected from the group consisting of  $\alpha$ -carboxy-2-nitrobenzyl; 1-(2-nitrophenyl)ethyl; 4,5-dimethoxy-2-nitrobenzyl; 1-(4,5-dimethoxy-2-nitrophenyl)ethyl; (4,5-dimethoxy-2-nitrobenzyloxy)carbonyl; 5-carboxymethoxy-2-nitrobenzyl; [(5-carboxymethoxy-2-nitrobenzyl)oxy]carbonyl; desoxybenzoinyl; and anthraquinon-2-ylmethoxycarbonyl.

25. (withdrawn) The method of Claim 16 wherein said molecule evidences switching based on electric (E) field induced band gap change, selected from the group consisting of:

(1) an E-field induced rotation of at least one rotatable section (rotor) of a molecule to change the band gap of the molecule (rotor/stator configuration);


(2) E-field-induced charge separation or recombination of the molecule via chemical bonding change to alter the band gap:

(2a) E-field-induced band gap change caused by the change of extended conjugation via charge separation or recombination accompanied by increasing or decreasing  $\pi$ - and/or p-electron localization;

(2b) E-field-induced band gap change caused by a change of extended conjugation via charge separation or recombination and  $\pi$ -bond breaking or formation; and

(3) E-field-induced band gap change via molecular folding or stretching.

26. (withdrawn) The method of Claim 25 wherein said bistable molecule comprises:



Example 1b

where:

A<sup>-</sup> is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

D<sup>+</sup> is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group is more electropositive than said Acceptor group;

Con<sub>1</sub> and Con<sub>2</sub> are connecting units between one molecule and another molecule or between a molecule and a substrate, said connecting units containing an attaching unit and at least one of said connecting units containing said photosensitive group, wherein said photosensitive group is selected from the group consisting of: photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive heteroring system(s) with at least one hetero-atom selected from the group consisting of N, O, S, B, and P and wherein the attaching unit is selected from the group consisting of carboxylic acid and its derivatives; sulfuric acid and its derivatives; phosphoric acid and its derivatives; hetero atoms selected from the group consisting of N, O, S, B, Se, and P; functional groups with at least one of said hetero atoms; hydrocarbons; and substituted hydrocarbons;

X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub> are tuning units built into the ring system which serve to tune the electronic properties, the optical properties, or both, of the bistable molecule as well as to that the ring system undergoes a smooth and desired tautomerization transition under the influence of an applied external E-field, wherein the tuning units are selected from the group consisting of a hetero atom selected from the group consisting of N, P, and As; hydrocarbons; and substituted hydrocarbons;


G<sub>1</sub> and G<sub>2</sub> are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more fragments to achieve desired molecular properties, wherein the bridging groups are either (a) selected from the group consisting of hetero atoms selected from the group consisting of N, O, S, and P; functional groups with at least one of

said hetero atoms; saturated or unsaturated hydrocarbons; and substituted hydrocarbons or (b) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

Q is a connecting unit between two phenyl rings, selected from the group consisting of S, O, NH, NR, hydrocarbons, and substituted hydrocarbons; and

H is a hydrogen atom.

27. (withdrawn) The method of Claim 25 wherein said bistable molecule comprises:



where:

A<sup>-</sup> is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

D<sup>+</sup> is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group is more electropositive than said Acceptor group;

Con<sub>2</sub> is a connecting unit between one molecule and another molecule or between a molecule and a substrate, said connecting unit containing an attaching unit and said photosensitive group, wherein said photosensitive group is selected from the group consisting of: photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive hetero-ring system(s) with at least one hetero-atom selected from the group consisting of N, O, S, B, and P and wherein the attaching unit is selected from the group consisting of carboxylic acid and its derivatives; sulfuric acid and its derivatives; phosphoric acid and its derivatives; hetero atoms selected from the group consisting of N, O, S, B, Se, and P; functional groups with at least one of said hetero atoms; hydrocarbons; and substituted hydrocarbons;

X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub> are tuning units built into the ring system which serve to tune the electronic properties, the optical properties, or both, of the bistable molecule as well as to ensure that the ring system undergoes a smooth and desired tautomerization transition under the influence of an applied external E-field, wherein the tuning units are selected from the group consisting of a hetero atom selected from the group consisting of N, P, and As, hydrocarbons, and substituted hydrocarbons;


G<sub>1</sub> and G<sub>2</sub> are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more fragments to achieve desired molecular properties, wherein the bridging groups are either (a) selected from the group consisting of hetero atoms selected from the group consisting of N, O, S, and P; functional groups with at least one of

said hetero atoms; saturated or unsaturated hydrocarbons; and substituted hydrocarbons or (b) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

Q is a connecting unit between two phenyl rings, selected from the group consisting of S, O, NH, NR, hydrocarbons, and substituted hydrocarbons; and

H is a hydrogen atom.

28. (withdrawn) The method of Claim 25 wherein said bistable molecule comprises:



where:

$A^-$  is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

$D^+$  is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group is more electropositive than said Acceptor group;


$G_1=G_2$ ,  $G_3=G_4$ ,  $G_5=G_6$ , and  $G_7=G_8$  are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more conjugated rings to achieve desired electronic properties, wherein the bridging groups are either (a) photosensitive functional groups or (b) selected from the group consisting of hetero atoms selected from the group consisting of N, O, S, and P; functional groups with at least one of said hetero atoms; saturated or unsaturated hydrocarbon; and substituted hydrocarbons, or (c) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

$Con_1$  and  $Con_2$  are connecting units between one molecule and another molecule or between a molecule and a substrate, said connecting units containing an attaching unit and at least one of said connecting units containing said photosensitive group, wherein said photosensitive group is selected from the group consisting of: photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive heteroring system(s) with at least one hetero-atom selected from the group consisting of N, O, S, B, and P and wherein the attaching unit is selected from the group consisting of carboxylic acid and its derivatives; sulfuric acid and its derivatives; phosphoric acid and its derivatives; hetero atoms selected from the group consisting of N, O, S, B, Se, and P; functional groups with at least one of said hetero atoms; hydrocarbons; and substituted hydrocarbons;

$R_1$ ,  $R_2$ , and  $R_3$  are spacing groups selected from the group consisting of (a) hydrogen, (b) saturated or unsaturated hydrocarbons, and (c) substituted hydrocarbons; and

$J_1$  and  $J_2$  are tuning groups to provide at least one appropriate functional effect selected from the group consisting of inductive effects, resonance effects, and steric effects; said tuning groups being selected from the group consisting of (a) hydrogen, (b) hetero atoms selected from the group consisting of N, O, S, P, B, F, Cl, Br, and I, (c) functional groups with at least one of said hetero atoms, (d) saturated or unsaturated hydrocarbons, and (e) substituted hydrocarbons.

29. (withdrawn) The method of Claim 25 wherein said bistable molecule comprises:



where:

$A^-$  is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;


$D^+$  is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated

hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group is more electropositive than said Acceptor group;

$R_1$ ,  $R_2$ , and  $R_3$  are spacing groups selected from the group consisting of (a) hydrogen, (b) saturated or unsaturated hydrocarbons, and (c) substituted hydrocarbons; and

$J_1$  and  $J_2$  are tuning groups to provide at least one appropriate functional effect selected from the group consisting of inductive effects, resonance effects, and steric effects; said tuning groups being selected from the group consisting of (a) hydrogen, (b) hetero atoms selected from the group consisting of N, O, S, P, B, F, Cl, Br, and I, (c) functional groups with at least one of said hetero atoms, (d) saturated or unsaturated hydrocarbons, and (e) substituted hydrocarbons.

30. (withdrawn) The method of Claim 25 wherein said bistable molecule comprises:



where:

$A^-$  is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of- (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one

of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

$D^+$  is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group is more electropositive than said Acceptor group;

$G_1=G_2$ ,  $G_3=G_4$ ,  $G_5=G_6$ , and  $G_7=G_8$  are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more conjugated rings to achieve desired electronic properties, wherein the bridging groups are either (a) photosensitive functional groups, or (b) selected from the group consisting of hetero atoms selected from the group consisting of N, O, S, and P; functional groups with at least one of said hetero atoms; saturated or unsaturated hydrocarbons; and substituted hydrocarbons, or (c) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

$Con_1$  and  $Con_2$  are connecting units between one molecule and another molecule or between a molecule and a substrate, said connecting units containing an attaching unit and at least one of said connecting units containing said photosensitive group, wherein said photosensitive group is selected from the group consisting of: photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive heteroring system(s) with at least one hetero-atom selected from the group consisting of N, O, S, B, and P and wherein the attaching unit is selected from the group consisting of carboxylic acid and its derivatives; sulfuric acid and its derivatives; phosphoric acid and its derivatives; hetero atoms selected from the group consisting of N, O, S, B, Se, and P; functional groups with at least one of said hetero atoms; hydrocarbons; and substituted hydrocarbons;

$J_1$ ,  $J_2$ ,  $J_3$ , and  $J_4$  are tuning groups which contain solvent functional groups selected from the group consisting of OH, NHR, COOH, and CN, where R is alkyl or aryl, wherein  $J_1$ -PSG,  $J_2$ -PSG,  $J_3$ -PSG, and  $J_4$ -PSG are linkages of said tuning groups with said photosensitive groups and are selected from the group consisting of ether, ester, carbonate, amide and carbamate linkages.

31. (new) A multiple electrode device comprising at least one pair of electrodes that form at least one junction and at least one connector species connecting said pair of electrodes in said junction, said junction having a functional dimension in nanometers or micrometers, wherein said at least one connector species comprises a bistable molecule provided  
5 with at least one photosensitive functional group for patterning said connector species.

32. (new) The multiple electrode device of Claim 31 wherein said at least one pair of electrodes comprises a positive terminal and a negative terminal and wherein said electrodes are adjacent each other.

33. (new) The multiple electrode device of Claim 32 wherein said second set of electrodes resides above said first set of electrodes, at a non-zero angle thereto.

34. (new) The multiple electrode device of Claim 32 wherein said second set of electrodes resides in the same plane as said first set of electrodes.

35. (new) The multiple electrode device of Claim 32 wherein said at least one pair of electrodes comprises said first set of electrodes and at least one probe addressing tip.

36. (new) The multiple electrode device of Claim 31 wherein said photosensitive functional group is sensitive to ultraviolet, electron-beam, or X-ray radiation.

37. (new) The multiple electrode device of Claim 31 wherein said bistable molecule comprises a main chain and at least one pendant group and wherein at least one photosensitive functional group is attached either to said main chain or to said pendant group.

38. (new) The multiple electrode device of Claim 37 wherein one said photosensitive group is attached to at least one end of said bistable molecule.

39. (new) The multiple electrode device of Claim 37 wherein said photosensitive group is selected from the group consisting of  $\alpha$ -carboxy-2-nitrobenzyl; 1-(2-nitrophenyl)ethyl; 4,5-dimethoxy-2-nitrobenzyl; 1-(4,5-dimethoxy-2-nitrophenyl)ethyl; (4,5-dimethoxy-2-

nitrobenzyloxy)carbonyl; 5-carboxymethoxy-2-nitrobenzyl; [(5-carboxymethoxy-2-nitrobenzyl)oxy]carbonyl; desoxybenzoinyl; and anthraquinon-2-ylmethoxycarbonyl.

40. (new) The multiple electrode device of Claim 31 wherein said molecule evidences switching based on electric (E) field induced band gap change, selected from the group consisting of:

(1) an electric field ("E-field") induced rotation of at least one rotatable section (rotor) of a molecule to change the band gap of the molecule (rotor/stator configuration);


(2) E-field-induced charge separation or recombination of the molecule via chemical bonding change to alter the band gap:

(2a) E-field-induced band gap change caused by the change of extended conjugation via charge separation or recombination accompanied by increasing or decreasing  $\pi$ - and/or p-electron localization;

(2b) E-field-induced band gap change caused by a change of extended conjugation via charge separation or recombination and  $\pi$ -bond breaking or formation; and

(3) E-field-induced band gap change via molecular folding or stretching.

41. (new) The multiple electrode device of Claim 40 wherein said bistable molecule comprises:



5 where:

$A^-$  is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its deriva-

tives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

$D^+$  is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH; (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P, wherein said Donor group is more electropositive than said Acceptor group;

$Con_1$  and  $Con_2$  are connecting units between one molecule and another molecule or between a molecule and a substrate, said connecting units containing an attaching unit and at least one of said connecting units containing said photosensitive group, wherein said photosensitive group is selected from the group consisting of photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive heteroring systems with at least one hetero-atom selected from the group consisting of N, O, S, B, and P,, and wherein the attaching unit is selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) hetero atoms selected from the group consisting of N, O, S, B, Se, and P, (e) functional groups with at least one of said hetero atoms (f) hydrocarbons, and (g) substituted hydrocarbons;

$X_1$ ,  $X_2$ ,  $X_3$  are tuning units built into the ring system which serve to tune the electronic properties, the optical properties, or both, of the bistable molecule as well those of the ring system undergo a smooth and desired tautomerization transition under the influence of an applied external E-field, wherein the tuning units are selected from the group consisting of a hetero atom selected from the group consisting of N, P, and As; hydrocarbons; and substituted hydrocarbons;


$G_1$  and  $G_2$  are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more fragments to achieve desired molecular properties, wherein the bridging groups are either (a) selected from the group consisting of (i) hetero atoms selected from the group consisting of N, O, S, and P; (ii) functional groups with at least one of said hetero atoms; (iii) saturated or unsaturated hydrocarbons; and (iv) substituted hy-

drocarbons or (b) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

Q is a connecting unit between two phenyl rings, selected from the group consisting of S, O, NH, NR, hydrocarbons, and substituted hydrocarbons; and

45 H is a hydrogen atom.

42. (new) The multiple electrode device of Claim 40 wherein said bistable molecule comprises:



5 where:

$A^-$  is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

$D^+$  is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P, wherein said Donor group is more electropositive than said Acceptor group;

$Con_2$  is a connecting unit between one molecule and another molecule or between a molecule and a substrate, said connecting unit containing an attaching unit and said photosensitive group, wherein said photosensitive group is selected from the group consisting of photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive hetero-ring systems with at least one hetero-atom selected from the group consisting of N, O, S, B, and P and wherein the attaching unit is selected from the group consisting of carboxylic acid and its derivatives; sulfuric acid and its derivatives; phosphoric acid and its derivatives; hetero atoms selected from the group consisting of N, O, S, B, Se, and P functional groups with at least one of said hetero atoms; hydrocarbons; and substituted hydrocarbons;

$X_1$ ,  $X_2$ ,  $X_3$  are tuning units built into the ring system which serve to tune the electronic properties, the optical properties, or both, of the bistable molecule as well as to that the ring system undergoes a smooth and desired tautomerization transition under the influence of an applied external E-field, wherein the tuning units are selected from the group consisting of a hetero atom selected from the group consisting of N, P, and As; hydrocarbons; and substituted hydrocarbons;

$G_1$  and  $G_2$  are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more fragments to achieve desired molecular properties, wherein the bridging groups are either (a) selected from the group consisting of (i) hetero atoms selected from the group consisting of N, O, S, and P; (ii) functional groups with at least

40

40

40

40



where:

5           A<sup>-</sup> is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

10           D<sup>+</sup> is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group is more electropositive than said Acceptor group;


15           G<sub>1</sub>=G<sub>2</sub>, G<sub>3</sub>=G<sub>4</sub>, G<sub>5</sub>=G<sub>6</sub>, and G<sub>7</sub>=G<sub>8</sub> are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more conjugated rings to achieve desired electronic properties, wherein the bridging groups are either (a) photosensitive functional groups or (b) selected from the group consisting of (i) hetero atoms selected from the group consisting of N, O, S, and P; (ii) functional groups with at least one of said hetero atoms; (iii) saturated or unsaturated hydrocarbons; and (iv) substituted hydrocarbons, or (c) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

20           Con<sub>1</sub> and Con<sub>2</sub> are connecting units between one molecule and another molecule or between a molecule and a substrate, said connecting units containing an attaching unit and at least one of said connecting units containing said photosensitive group, wherein said photosensitive group is selected from the group consisting of photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive hetero-ring systems with at least one hetero-atom selected from the group consisting of N, O, S, B, and P and wherein the attaching unit is selected from the group consisting of carboxylic acid and its derivatives; sulfuric acid and its derivatives; phosphoric acid and its derivatives; hetero atoms selected from the group consisting of N, O, S, B, Se, and P; functional groups with at least one of said hetero atoms; hydrocarbons, and substituted hydrocarbons;

$R_1$ ,  $R_2$ , and  $R_3$  are spacing groups selected from the group consisting of (a) hydrogen, (b) saturated or unsaturated hydrocarbons, and (c) substituted hydrocarbons; and

$J_1$  and  $J_2$  are tuning groups to provide at least one appropriate functional effect selected from the group consisting of inductive effects, resonance effects, and steric effects, where said tuning groups are selected from the group consisting of (a) hydrogen, (b) hetero atoms selected from the group consisting of N, O, S, P, B, F, Cl, Br, and I, (c) functional groups with at least one of said hetero atoms, (d) saturated or unsaturated hydrocarbons, and (e) substituted hydrocarbons.

44. (new) The multiple electrode device of Claim 40 wherein said bistable molecule comprises:



5 where:


$A^-$  is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its derivatives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

$D^+$  is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group is more electropositive than said Acceptor group;

$R_1$ ,  $R_2$ , and  $R_3$  are spacing groups selected from the group consisting of (a) hydrogen, (b) saturated or unsaturated hydrocarbons, and (c) substituted hydrocarbons; and

$J_1$  and  $J_2$  are tuning groups to provide at least one appropriate functional effect selected from the group consisting of inductive effects, resonance effects, and steric effects, said tuning groups are selected from the group consisting of (a) hydrogen, (b) hetero atoms selected from the group consisting of N, O, S, P, B, F, Cl, Br, and I, (c) functional groups with at least one of said hetero atoms, (d) saturated or unsaturated hydrocarbons, and (e) substituted hydrocarbons.

45. (new) The multiple electrode device of Claim 40 wherein said bistable molecule comprises:



5 where:

$A^-$  is an Acceptor group comprising an electron-withdrawing group selected from the group consisting of (a) carboxylic acid and its derivatives, (b) sulfuric acid and its deriva-

tives, (c) phosphoric acid and its derivatives, (d) nitro, (e) nitrile, (f) hetero atoms selected from the group consisting of N, O, S, P, F, Cl, and Br, (g) functional groups with at least one of said hetero atoms, (h) saturated or unsaturated hydrocarbons, and (i) substituted hydrocarbons;

$D^+$  is a Donor group comprising an electron-donating group selected from the group consisting of (a) hydrogen, (b) amines, (c) OH, (d) SH, (e) ethers, (f) saturated or unsaturated hydrocarbons, (g) substituted hydrocarbons, and (h) functional groups with at least one hetero atom selected from the group consisting of B, Si, I, N, O, S, and P; wherein said Donor group is more electropositive than said Acceptor group;

$G_1=G_2$ ,  $G_3=G_4$ ,  $G_5=G_6$ , and  $G_7=G_8$  are bridging groups for connecting stator and rotor portions of said bistable molecule or for connecting two or more conjugated rings to achieve desired electronic properties, wherein the bridging groups are either (a) photosensitive functional groups, or (b) selected from the group consisting of (i) hetero atoms selected from the group consisting of N, O, S, and P; (ii) functional groups with at least one of said hetero atoms; (iii) saturated or unsaturated hydrocarbons; and (iv) substituted hydrocarbons, or (c) selected from the group consisting of a single atom bridge and a direct sigma bond between said rotor and stator portions;

$Con_1$  and  $Con_2$  are connecting units between one molecule and another molecule or between a molecule and a substrate, said connecting units containing an attaching unit and at least one of said connecting units containing said photosensitive group, wherein said photosensitive group is selected from the group consisting of photosensitive azo, photosensitive ester, photosensitive ether, photosensitive amide, photosensitive imide, photosensitive amine, photosensitive imine, photosensitive carbonate, photosensitive carbamate, photosensitive thio-ether, photosensitive thio-ester, photosensitive isocyanides, and photosensitive heteroring systems with at least one hetero-atom selected from the group consisting of N, O, S, B, and P and wherein the attaching unit is selected from the group consisting of carboxylic acid and its derivatives; sulfuric acid and its derivatives; phosphoric acid and its derivatives; hetero atoms selected from the group consisting of N, O, S, B, Se, and P; functional groups with at least one of said hetero atoms; hydrocarbons; and substituted hydrocarbons; and

$J_1$ ,  $J_2$ ,  $J_3$ , and  $J_4$  are tuning groups which contain solvent functional groups selected from the group consisting of OH, NHR, COOH, and CN, where R is alkyl or aryl, wherein  $J_1$ -PSG,  $J_2$ -PSG,  $J_3$ -PSG, and  $J_4$ -PSG are linkages of said tuning groups with said photosensitive

40 groups and are selected from the group consisting of ether, ester, carbonate, amide, and carbamate linkages.